

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

# (12) UK Patent Application (19) GB (11) 2 270 312 (13) A

(43) Date of A Publication 09.03.1994

(21) Application No 9317770.7

(22) Date of Filing 26.08.1993

(30) Priority Data

(31) 939258

(32) 02.09.1992

(33) US

(71) Applicant(s)

Merck & Co Inc

(Incorporated in USA - New Jersey)

P O Box 2000, 126 East Lincoln Avenue, Rahway,  
New Jersey 07065-0900, United States of America

(72) Inventor(s)

Claude Dufresne  
Prakash S Masurekar  
Leslie Anne Ferrell  
Deborah L Zink  
Margaret S Sosa

(74) continued overleaf

(51) INT CL<sup>5</sup>

C07D 493/08, A61K 31/34 // (C07D 493/08 307:24  
319:06)

(52) UK CL (Edition M)

C2C CAA CBC C1485 C200 C214 C22Y C220 C226 C25Y  
C253 C29X C29Y C292 C30Y C306 C36Y C360 C362  
C363 C366 C367 C368 C491 C633 C643 C648 C65X  
C652 C658 C67X C672 C678 C699 C80Y C800 C801  
C802  
U1S S1313 S2410 S2415

(56) Documents Cited


EP 0475706 A US 5026554 A

(58) Field of Search


UK CL (Edition L) C2C CBC  
INT CL<sup>5</sup> C07D  
ONLINE DATABASES: CAS ONLINE

(54) Cholesterol lowering compounds


(57) New Zaragozic acids have been isolated from a culture of MF5465. These compounds and their derivatives are active as squalene synthetase inhibitors and are useful in the treatment of hypercholesterolemia or cancer or as antifungal agents. Included are compounds of the formula:-



wherein R¹ is



R² is



with certain exceptions, and Z¹, Z² and Z³ are each hydrogen or certain defined esterifying groups.

GB 2 270 312 A

AP2

**(74) Agent and/or Address for Service**

**H K Quillin**

**Merck & Co Inc, European Patent Department,  
Terlings Park, Eastwick Road, HARLOW, Essex,  
CM20 2QR, United Kingdom**

2270312

5

- 1 -

10 TITLE OF THE INVENTION  
CHOLESTEROL LOWERING COMPOUNDS

BACKGROUND OF THE INVENTION

15 Hypercholesterolemia is known to be one of  
the prime risk factors for ischemic cardiovascular  
disease, such as arteriosclerosis. Bile acid  
sequestrants have been used to treat this condition;  
they seem to be moderately effective but they must be  
20 consumed in large quantities, i.e. several grams at a  
time, and they are not very palatable.

MEVACOR® (lovastatin), now commercially  
available, is one of a group of very active  
antihypercholesterolemic agents that function by  
25 limiting cholesterol biosynthesis by inhibiting the  
enzyme, HMG-CoA reductase.

Squalene synthetase is the enzyme involved in  
the first committed step of the de novo cholesterol  
biosynthetic pathway. This enzyme catalyzes the  
30 reductive dimerization of two molecules of farnesyl

pyrophosphate to form squalene. The inhibition of this committed step to cholesterol should leave unhindered biosynthetic pathways to ubiquinone, dolichol and isopentenyl t-RNA.


5 Previous efforts at inhibiting squalene synthetase have employed pyrophosphate or pyrophosphate analog containing compounds such as those described in P. Ortiz de Montellano et al, J. Med Chem. 20, 243 (1977) and E.J. Corey and R. Volante, J. Am. Chem. Soc., 98, 1291 (1976). S. Biller (U.S. Patent 10 4,871,721) describes isoprenoid (phosphinylmethyl)phosphonates as inhibitors of squalene synthetase.

15 Recently certain nonphosphorous containing inhibitors of squalene synthetase have been isolated as natural products. These natural product inhibitors are described in U.S. Patents 5,053,425; 5,055,487 and 5,026,554. Certain squalenestatsins are discussed in Baxter, A. et al., J. Biol. Chem. 267, 11705 (1992); 20 Dawson, M.J. et al., J. Antib. 45, 639 (1992); Sidebottom, P.J. et al., J. Antib. 45, 648 (1992). See also EPO 494622 and PCT WO 92/12156.

U.S. Patent 5,026,554 describes a squalene synthetase inhibitor of structure:

25


30



which is hereafter referred to as Zaragozic acid C. This compound is produced by a solid fermentation employing a culture of Leptodontium elatius. The present invention discloses certain structurally related compounds to Zaragozic acid C which have now been isolated as minors from a fermentation broth using Leptodontium elatius.



#### DETAILED DESCRIPTION OF THE INVENTION


The compounds of the structural formula (I), or their pharmaceutically acceptable salts, are useful in the treatment of hypercholesterolemia, fungal infections, and cancer. Structural formula (I) consists of




or pharmaceutically acceptable salt thereof, wherein

$R^1$  is selected from the group consisting of







5  $R^2$  is selected from the group consisting of




10



15



20



25  $z^1$ ,  $z^2$  and  $z^3$  are each independently selected from;

a) H;

b)  $C_{1-5}$ alkyl;

c)  $C_{1-5}$ alkyl substituted with


a member of the group consisting of:

i) phenyl,

30


ii) phenyl substituted with methyl,  
methoxy, halogen (Cl, Br, I, F) or  
hydroxy;

with the proviso that when  $R^2$  is




5

then  $R^1$  is neither



10

nor,



15




One embodiment of the present invention is  
20 formula I compounds further restricted to compounds  
wherein  $Z^1$ ,  $Z^2$  and  $Z^3$  are each independently H or CH<sub>3</sub>.

Another embodiment is a compound selected  
from the group consisting of the following structures,

25





30





25

30



or a pharmaceutically acceptable salt thereof.

The present invention is directed to certain zaragozic acid derivatives isolated from the fermentation of MF 5465.

5 The compounds of formula (I) can be prepared in an aerobic fermentation procedure employing a fungal culture, MF5465, identified as Leptodontium elatius. This fermentation procedure is described in U.S. Patent 5,026,554, the contents of which are herein specifically incorporated by reference.

10 The compounds of the present invention have been isolated as components produced in lower yield than Zaragozic acid C from this fermentation procedure.

15 Mutants of MF5465, having essentially the same characteristics as MF 5465, are also capable of producing compounds of this invention.

The culture MF5465 is that of a fungus, a lignicolous Hyphomycete, Leptodontium elatius, isolated from wood in the Joyce Kilmer Memorial Forest in North Carolina. This culture has been deposited with the  
20 American Type Culture Collection at 12301 Parklawn Drive, Rockville, MD 20852 as ATCC 74011.

The characteristics of MF5465 are described in U.S. Patent 5,026,554.

25 Compounds of this invention can be obtained by culturing the above noted microorganism in an aqueous nutrient medium containing sources of assimilable carbon and nitrogen, preferably under aerobic conditions. Nutrient media may also contain mineral salts and defoaming agents.

30 The preferred sources of carbon in the nutrient medium are carbohydrates such as glycerin, glucose, starch, dextrin, and the like. Other sources which may be included are maltose, mannose, sucrose,

and the like. In addition, complex nutrient sources such as oat flour, corn meal, millet, corn and the like may supply utilizable carbon. The exact quantity of the carbon source which is used in the medium will  
5 depend, in part, upon the other ingredients in the medium, but is usually found in an amount ranging between 0.5 and 5 percent by weight. These carbon sources can be used individually in a given medium or several sources in combination in the same medium.

10 The preferred sources of nitrogen include complex sources such as peptone, yeast extracts (hydrolysates, autolysates), dried yeast, tomato paste, soybean meal, corn steep liquor, distillers solubles, malt extracts and the like. Inorganic nitrogen sources  
15 such as ammonium salts (eg. ammonium nitrate, ammonium sulfate, ammonium phosphate, etc.) or other nitrates or nitrites can also be used. The various sources of nitrogen can be used alone or in combination in amounts ranging between 0.2 to 70 percent by weight of the  
20 medium.

The carbon and nitrogen sources are generally employed in combination, but need not be in pure form. Less pure materials which contain traces of growth factors, vitamins, and mineral nutrients may also be  
25 used. Mineral salts may also be added to the medium such as (but not limited to) calcium carbonate, sodium or potassium phosphate, sodium or potassium chloride, magnesium salts, copper salts, cobalt salts and the like. Also included are trace metals such as  
30 manganese, iron, molybdenum, zinc, and the like. In addition, if necessary, a defoaming agent such as polypropylene glycol or silicone may be added, especially if the culture medium foams seriously.

5       The preferred process for production of compounds of this invention consists of inoculating spores or mycelia of the producing organism into a suitable medium and then cultivating under aerobic condition.

10       The fermentation procedure begins by inoculating a preserved source of culture (spores or mycelia) into a nutrient seed medium and obtaining, sometimes through a multi step process, growth of the organism which serves as inoculum for the production of the active compounds. After inoculation, the flasks are incubated with rotary shaking at temperatures ranging from 20 to 30°C, preferably 25 to 28°C. Shaking rates may range up to 400 rpm, preferably 200 to 220 rpm. Seed flasks are incubated over a period of 2 to 10 days, preferably 2 to 4 days. When biomass is plentiful, usually 2 to 4 days, the culture may be used to inoculate production medium flasks. Additional seed stages are required when using larger vessels for production. When this is done, a portion of the culture is used to inoculate a subsequent seed flask or fermentor which is incubated under similar conditions.

20       After inoculation, the production fermentation is incubated for 3 to 30 days, preferably 7 to 21 days, with or without agitation (depending on whether liquid or solid fermentation media are employed). The fermentation is conducted aerobically at temperatures ranging from 20 to 40°C. If used, agitation or shaking may be at a rate of 100 to 400 rpm. To obtain optimum results, the temperatures are in the range of 22 to 28°C, most preferably 24 to 26°C. The pH of the nutrient medium suitable for producing the active compounds is in the range of 3.5

to 8.5, most preferably 5.0 to 7.5. After the appropriate period for production of the desired compound, fermentation flasks or fermentors are harvested and the active compound isolated.

5           An organic solvent is employed to extract a compound of this invention from the fermentation medium. The preferred solvent for extraction of the fermentation broth is a water immiscible solvent such as isopropyl acetate. The mixture of organic solvent  
10           and fermentation broth is vigorously stirred and filtered. The organic extract is then adsorbed on an anion exchange resin. The preferred resin is a weakly basic anion exchange resin such as AMBERLYST A-21 (acetate cycle). The active compounds can be eluted  
15           from AMBERLYST A-21 using a low pH solution or a high salt eluant. The preferred eluant is 3% ammonium chloride in 90% methanol/water. After elution from the ion exchange resin, the active compounds may be  
20           recovered from the eluate by diluting the eluate with water, acidifying to  $\text{pH} < 3$ , and extracting into an organic solvent. The preferred solvent for extraction is ethyl acetate. The organic extract is then evaporated to afford a partially purified complex of active compounds.

25           The complex of active compounds is further purified by chromatographic separation which may be carried out by employing reverse phase chromatography. The HPLC column packing, also known as adsorbent, may include octyl, octadecyl, phenyl, cyano, or diol bonded  
30           silica gel, alumina, or polymer. The preferred adsorbent for this chromatography is a C8 bonded phase silica gel. The mobile phase of the eluant comprises methanol, acetonitrile, or tetrahydrofuran as an organic

modifier for an aqueous solution buffered at a pH in the range of 2 to 8. The preferred eluant for reverse phase chromatography is a mixture of acetonitrile and water buffered at a low pH, such as 0.1% phosphoric acid, or trifluoroacetic acid. The flow rate ranges from 0.5 to 4 ml/min for a column of approximately 5 mm ID. The temperature of the column ranges between 10°C and 45°C.

The relative retention time (RRT) is the ratio of the retention time of a given major peak under given chromatographic conditions, divided by the retention time of Zaragozic acid C under identical chromatographic conditions. The structure of Zaragozic acid C is provided in the Background of the Invention, above.

The present invention is also directed to a method of treating hypercholesterolemia which comprises the administration to a subject in need of such treatment a nontoxic therapeutically effective amount of a compound represented by structural formula (I) and pharmaceutically acceptable salts thereof. Specifically, the compounds of this invention are useful as antihypercholesterolemic agents for the treatment of arteriosclerosis, hyperlipidemia, familial hypercholesterolemia and the like diseases in humans. They may be administered orally or parenterally in the form of a capsule, a tablet, an injectable preparation or the like. It is usually desirable to use the oral route. Doses may be varied, depending on the age, severity, body weight and other conditions of human patients, but daily dosage for adults is within a range of from about 20 mg to 2000 mg (preferably 20 to 100

mg) which may be given in two to four divided doses. Higher doses may be favorably employed as required.

The present invention is also directed to a method of inhibiting squalene synthetase which  
5 comprises the administration to a subject in need of such treatment a nontoxic therapeutically effective amount of a compound represented by structural formula (I) and pharmaceutically acceptable salts thereof. . Specifically, the compounds of this invention are  
10 useful in treating disease conditions such as, but not limited to, hypercholesterolemia which result from the action of the enzyme squalene synthetase. They may be administered orally or parenterally in the form of a capsule, a tablet, an injectable preparation or the  
15 like. It is usually desirable to use the oral route. Doses may be varied, depending on the age, severity, body weight and other conditions of human patients, but daily dosage for adults is within a range of from about 20 mg to 2000 mg (preferably 20 to 100 mg) which may be  
20 given in two to four divided doses. Higher doses may be favorably employed as required.

The pharmaceutically acceptable salts of the compounds of this invention include those formed from cations such as sodium, potassium, aluminum, calcium,  
25 lithium, magnesium, zinc, and from bases such as ammonia, ethylenediamine, N-methyl-glutamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylene-diamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, diethylamine, piperazine,  
30 tris(hydroxymethyl)aminomethane, and tetramethylammonium hydroxide. The salts included herein encompass those wherein one, two or all three of the carboxyl groups are in the salt form.



The compounds of this invention may also be administered in combination with other cholesterol lowering agents such as those which inhibit an enzymatic pathway in the biosynthesis of cholesterol. Example of such agents would include but are not limited to HMG-CoA reductase inhibitors, HMG-CoA synthase inhibitors, and squalene epoxidase inhibitors. Illustrative of such inhibitors are lovastatin, simvastatin, pravastatin and fluvastatin. Other cholesterol lowering agents that may be administered include niacin, probucol, and the fibric acids, clofibrate and gemfibrozil. Appropriate daily dosages for adults are niacin (2-8 gm), probucol (up to 1000 mg), clofibrate (up to 2 gm) and gemfibrozil (800-1500 mg).

The compounds of this invention may also be coadministered with pharmaceutically acceptable nontoxic cationic polymers capable of binding bile acids in a non-reabsorbable form in the gastrointestinal tract. Examples of such polymers include cholestyramine, colestipol and poly[methyl-(3-trimethylaminopropyl)imino-trimethylene dihalide]. The relative amounts of the compounds of this invention and these polymers is between 1:100 and 1:15,000.

The intrinsic squalene synthetase inhibitory activity of representative compounds of this invention is measured by the standard in vitro protocol described below:

#### Preparation of Microsomes:

Male, Charles River CD rats (120 to 150 g) are fed a diet containing 0.1% lovastatin for 4 days. The livers from these rats were homogenized in 5

volumes (ml/g) of ice cold 50 mM HEPES (4-(2- hydroxy-ethyl)-1-piperazine-ethanesulfonic acid), 5 mM EDTA(ethylenediaminetetraacetic acid) pH 7.5 with a Potter-Elvehjem type tissue grinder. The homogenate is  
5 centrifuged twice at 20,000 x g for 15 minutes at 4°C, discarding the pellet each time. The supernatant is then centrifuged at 100,000 x g for 1 hour at 4°C. The resulting microsomal pellet is resuspended in a  
10 volume of the above homogenizing buffer equal to one-fifth the volume of the original homogenate. This microsomal preparation typically has a protein concentration of about 7 mg/ml. The microsomal suspensions are stored in aliquots at -70°C. Squalene synthetase activity in these aliquots is stable for at  
15 least several months.

#### Partial Purification of Prenyl Transferase

Prenyl transferase is purified to use in the  
20 enzymatic synthesis of radiolabelled farnesyl pyrophosphate. Prenyl transferase is assayed by the method of Rilling (Methods in Enzymology 110, 125-129 (1985)) and a unit of activity is defined as the amount of enzyme that will produce 1 μmole of farnesyl  
25 pyrophosphate per minute at 30°C in the standard assay.

The livers of 23 forty-day old male rats that have been fed 5% cholestyramine plus 0.1% lovastatin are homogenized in a Waring blender in 1 liter of 10 mM mercaptoethanol, 2 mM EDTA, 25 μM leupeptin, 0.005%  
30 phenylmethyl sulfonyl fluoride pH 7.0 containing 0.1 trypsin inhibitor units of aprotinin/ml. The homogenate is centrifuged at 20,000 x g for 20

minutes. The supernatant is adjusted to pH 5.5. with 6N HOAc and centrifuged at 100,000 x g for 1 hour. This supernatant is adjusted to pH 7.0 with 3N KOH and a 35-60% ammonium sulfate fraction taken. The 60%  
5 pellet is redissolved in 60 ml of 10 mM potassium phosphate, 10 mM mercaptoethanol, 1 mM EDTA pH 7.0 (Buffer A) and dialyzed against two 1 liter changes of Buffer A. This dialyzed fraction is applied to a 12.5 x 5 cm column of DEAE-sepharose 4B equilibrated with  
10 Buffer A. The column is washed with 700 ml of Buffer A and a 1 liter gradient from Buffer A to 100 mM potassium phosphate, 10 mM mercaptoethanol, 1 mM EDTA pH 7.0. Fractions having a specific activity greater than 0.20 units/mg are combined, solid ammonium sulfate  
15 is added to bring to 60% saturation and pelleted. The pellet is dissolved in 8 ml of 10 mM Tris, 10 mM  $\beta$ -mercaptoethanol pH 7.0 (Buffer B). The redissolved pellet is taken to 60% saturation with ammonium sulfate by adding 1.5 volumes of saturated ammonium sulfate in  
20 Buffer B. This ammonium sulfate suspension typically contains 3.5 units/ml with a specific activity of 0.23 units/mg and is free of isopentenyl pyrophosphate isomerase activity. This ammonium sulfate suspension is used for the synthesis of [4- $^{14}$ C]farnesyl-  
25 pyrophosphate and its activity is stable stored at 4°C for at least 6 months.

#### Enzymatic Synthesis of [4- $^{14}$ C]farnesyl-pyrophosphate

30 The solvent (ethanol: 0.15 N  $\text{NH}_4\text{OH}$ , 1:1) is removed from 55  $\mu\text{Ci}$  of [4- $^{14}$ C]isopentenyl pyrophosphate (47.9  $\mu\text{Ci}/\mu\text{mole}$ ) by rotary evaporation. Six

hundred microliters of 100 mM Tris, 10 mM  $\text{MgCl}_2$ , 4 mM dithiothreitol pH 7.5 is added and the solution is transferred to a 1.5 ml Eppendorf centrifuge tube. Geranyl-pyrophosphate, 250  $\mu\text{l}$  of a 20 mM solution, and  
5 50  $\mu\text{l}$  of the ammonium sulfate suspension of prenyl transferase are added to initiate the reaction. This incubation contains 5  $\mu\text{moles}$  of geranyl pyrophosphate, 1.15  $\mu\text{moles}$  of isopentenyl pyrophosphate, 6  $\mu\text{moles}$  of  $\text{MgCl}_2$  of 0.18 units of prenyl transferase in a volume  
10 of 900  $\mu\text{l}$ . The incubation is conducted at 37°C. During the incubation, the mix typically turns cloudy white as the newly formed magnesium complex of farnesyl pyrophosphate precipitates out of solution. The  
[4- $^{14}\text{C}$ ]farnesyl pyrophosphate is collected by  
15 centrifugation for 3 minutes at 14,000 rpm in an centrifuge tube, the supernatant removed, and the pellet is dissolved in 1.0 ml of 50 mM HEPES, 5 mM EDTA, pH 7.5. The yield is typically about 50  $\mu\text{Ci}$  (90%)  
of [4- $^{14}\text{C}$ ]farnesyl pyrophosphate. The [4- $^{14}\text{C}$ ]farnesyl  
20 pyrophosphate is stored in aliquots at -70°C.

#### Squalene Synthetase Assay

Reactions are performed in 16 x 125 mm screw  
25 cap test tubes. A batch assay mix is prepared from the following solution:

30

		<u>ml per</u> <u>assay</u>	<u>volume for</u> <u>50 assays</u>
	1. 250 mM Hepes pH 7.5	20	1000
	2. NaF 110 mM	10	500
5.	3. MgCl <sub>2</sub> 55 mM	10	500
	4. Dithiothreitol 30 mM	10	500
	5. NADPH 10 mM (made fresh)	10	500
	6. [4- <sup>14</sup> C]farnesyl-pyrophosphate		
	47.9 $\mu$ Ci/ $\mu$ mole, and	3.0	150
10	0.025 $\mu$ Ci/3.0 $\mu$ l		
	7. H <sub>2</sub> O	24	1200

This assay mix is degassed under a vacuum and  
 flushed with N<sub>2</sub>. Solutions of the squalene synthetase  
 15 inhibitors are prepared either in DMSO or MeOH and a  
 1:120 dilution of the microsomal protein is made with  
 the original homogenizing buffer. For each reaction,  
 87  $\mu$ l of the assay mix is taken with 3  $\mu$ l of an  
 inhibitor solution (DMSO or MeOH in the controls),  
 20 warmed to 30°C in a water bath and then the reaction is  
 initiated by the addition of 10  $\mu$ l of the 1:120  
 dilution of microsomal protein (0.6  $\mu$ g protein total in  
 the assay). The reactions are stopped after 20 minutes  
 by the addition of 100  $\mu$ l of a 1:1 mix of 40% KOH with  
 25 95% EtOH. The stopped mix is heated at 65°C for 30  
 minutes, cooled, 10 ml of heptane is added and the mix  
 is vortexed. Two grams of activated alumina is then  
 added, the mix vortexed again, the alumina allowed to  
 settle and 5 ml of the heptane layer is removed. Ten  
 30 ml of scintillation fluid is added to the heptane  
 solution and radioactivity is determined by liquid  
 scintillation counting.

Percent inhibition is calculated by the formula:

$$1 - \frac{[\text{Sample} - \text{Blank}]}{[\text{Control} - \text{Blank}]} \times 100$$

5

IC<sub>50</sub> values are determined by plotting the log of the concentration of the test compound versus the percentage inhibition. The IC<sub>50</sub> is the concentration of inhibitor that gives 50% inhibition as determined from these plots.

The present compounds are also useful as broad spectrum antifungal agents as determined by broth and agar dilution methods. Thus the present invention is also directed to a method of treating fungus infections which comprises the administration to an organism in need of such treatment a nontoxic therapeutically effective amount of a compound represented by the structural formula (I) and pharmaceutically acceptable salts thereof. Generally from 2 to about 20 mg/kg should be employed as a unit dosage in an antifungal treatment.

Furthermore the compounds of the present invention are useful as inhibitors of farnesyl-protein transferase and thereby of farnesylation of the RAS protein and thus block the ability of RAS to transform normal cells to cancer cells. Farnesyl-protein transferase activity may be reduced or completely inhibited by adjusting the compound dose.

30

The intrinsic farnesyl-protein transferase (Ftase) activity of representative compounds of this invention is measured by the assays described below:

5

#### RASIT ASSAY I

Farnesyl-protein transferase (Ftase) from bovine brain is chromatographed on DEAE-Sephacel (Pharmacia, 0-0.8 M NaCl gradient elution), N-octyl  
10 agarose (Sigma, 0-0.6 M NaCl gradient elution), and a mono Q HPLC column (Pharmacia, 0-0.3 M NaCl gradient). Ras-CVLS at 3.5  $\mu$ M, 0.25  $\mu$ M [ $^3$ H]FPP, and the compounds to be assayed are incubated with this partially  
15 purified enzyme preparation.

15

#### RASIT ASSAY II

Farnesyl-protein transferase (Ftase) from bovine brain is chromatographed on DEAE-Sephacel  
20 (Pharmacia, 0-0.8 M NaCl gradient elution), N-octyl agarose (Sigma, 0-0.6 M NaCl gradient elution), and a mono Q HPLC column (Pharmacia, 0-0.3 M NaCl gradient). Ras-CVLS at 1.0  $\mu$ M, 0.5  $\mu$ M [ $^3$ H]FPP, and the compounds to be assayed are incubated with this partially  
25 purified enzyme preparation. The Ftase data is a measurement of the ability of the test compound to inhibit Ras farnesylation in vitro.

The pharmaceutical compositions containing the compounds of structural formula I inhibit  
30 farnesyl-protein transferase and the farnesylation of the oncogene protein Ras. These compounds are useful as pharmaceutical agents for mammals, especially for

humans. These compounds may be administered to patients for use in the treatment of cancer. Examples of the type of cancer which may be treated with the compounds of this invention include, but are not  
5 limited to, colorectal carcinoma, exocrine pancreatic carcinoma, and myeloid leukemias.

The present invention also encompasses a method of the treatment of cancer, comprising the administration of a pharmaceutical composition  
10 comprising a therapeutically effective amount of the compounds of this invention, with or without pharmaceutically acceptable carriers or diluents.

Suitable compositions of this invention include aqueous solutions comprising compounds of this  
15 invention and pharmacologically acceptable carriers, e.g. saline, at a pH level, e.g., 7.4. The solutions may be introduced into a patient's intramuscular blood-stream by local bolus injection.

When a compound according to this invention  
20 is administered into a human subject, the daily dosage will normally be determined by the prescribing physician with the dosage generally varying according to the age, weight, and response of the individual patient, as well as the severity of the patient's  
25 symptoms.

In one exemplary application, a suitable amount of compound is administered to a human patient undergoing treatment for cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to  
30 about 20 mg/kg of body weight of a mammal per day, preferably of between 0.5 mg/kg of body weight to about 10 mg/kg of body weight of a mammal a day.



The following Examples illustrate the preparation of the compounds of formula (I) and their incorporation into pharmaceutical compositions and, as such, are not to be considered as limiting the invention set forth in the claims appended hereto.

#### EXAMPLE 1

##### Fermentation of *Leptodontium elatius* at the 500 L Scale

###### Seed Stock Preparation

*Leptodontium elatius* (MF5465) was obtained from the MSDRL culture collection as a lyophilized preparation (order #, N920316001). The working seed stock was prepared by inoculating a 250 ml flask containing 50 ml of seed medium with the lyophilized culture and incubating for 48 hours at 25°C with 220 RPM rotary shaking. The seed medium contained per liter in deionized water: soluble starch, 30 gm; cerelose, 10 gm; pharmamedia, 10 gm;  $\text{KH}_2\text{PO}_4$ , 9 gm; DIFCO yeast extract (8005), 5 gm;  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.2 gm;  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 10 mg;  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ , 10 mg;  $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ , 0.25 mg;  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 0.75;  $\text{H}_3\text{BO}_4$ , 0.56 mg;  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , 2.0 mg;  $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ , 0.19 mg, and the pH was adjusted to 6.0 with sodium hydroxide. After the 48 hour incubation period, 2 ml of the culture were transferred to each of a second set of 4 four flasks containing seed medium as described above. Incubation conditions were identical to those described above. After incubation of the 4 second generation cultures, the 200 ml of culture broth contained in the 4 flasks was combined with 100 ml of 30% glycerol and the resultant mixture was aliquoted at 1 ml/cryovial and stored at

-70°C. These aliquots of frozen vegetative mycelia (FVM) were used to initiate inoculum development.

#### Inoculum Development

5           One FVM was used to inoculate each 250 ml flask containing 50 ml of seed medium. The cultures were incubated at 25°C with 220 RPM rotary shaking for 48 hours. Twenty milliliters were transferred to each  
10       2 L flask containing 500 ml of seed medium. The cultures were incubated at 25°C with 220 RPM rotary shaking for 48 hours. Four 2 L scale cultures were combined (2 L total culture volume) and used to inoculate a 285 L fermentor containing 180 L of seed  
15       medium (with 2 ml/L polypropylene glycol to suppress foaming). The fermentation was carried out for 48 hours under the following conditions: temperature, 25°C; pressure, 0.7 Kg/cm<sup>2</sup>; air flow, 75 liters/min.; agitation, 200 RPM.

#### 20       Production Fermentation

          Twenty-five liters from the seed fermentation were inoculated into a 760 liter fermentor containing 500 L of production medium which contained per liter in deionized water: primatone HS, 23 gm; DIFCO yeast  
25       extract (8005), 7.5 gm; sodium citrate, 11 gm; glycerol (sterilized separately as a 50% w/v solution), 20 gm; polypropylene glycol, 0.5 ml, and the pH was adjusted to 6.0 with sulfuric acid. The medium was sterilized for 30 minutes at 123°C and the fermentation was  
30       carried out for approximately 19 days under the following conditions: temperature, 25°C; pressure, 0.7 Kg/cm<sup>2</sup>; air flow, 100 liters/min.; agitation, 160 RPM. pH was allowed to rise from approximately 6.0 (post-

inoculation) and was controlled at 7.2 with sulfuric acid. Residual glycerol was monitored and as the concentration approached 5 gm/L, glycerol was added to the fermentor as a sterile 50% w/v solution to give a  
5. final concentration of approximately 20 gm/L.

#### EXAMPLE 2

##### Isolation Process:

The whole broth (1500 L, which is three  
10 combined batches prepared by Example 1) was acidified to pH 2.3 with sulfuric acid. The broth was extracted twice with isopropyl acetate (1500 L each time). (For each extraction, isopropyl acetate was added and the mixture was agitated for 1 hour. The isopropyl acetate  
15 extract was recovered from the mixture using a Westfalia Decanter.) The two extracts were combined (3000 L) and applied to an AMBERLYST A-21 (OAc cycle) column (132 L bed volume). The resin was prepared by stirring it in 1N acetic acid before packing into the  
20 column. The resin bed was successively washed with MeOH (132 L) and isopropyl acetate (132 L). After the isopropyl acetate extract was passed through the column, the column was washed with 50% MeOH/H<sub>2</sub>O (132 L). The spent and wash were discarded. The column was  
25 then eluted with 3% NH<sub>4</sub>Cl in 90% MeOH/H<sub>2</sub>O (1320 L). The eluate (1320 L) was diluted with water (402 L) and loaded onto an HP-20 column (95 L). The spent and wash were discarded. The column was eluted with 90% MeOH/H<sub>2</sub>O (380 L). The HP-20 eluate was diluted with  
30 water (380 L) and adjusted to pH 2.3 with concentrated sulfuric acid. EtOAc (570 L) and water (380 L) were then added. The EtOAc layer (492 L) was separated and concentrated under reduced pressure to an oily residue.

The residue was dissolved in acetonitrile (500 mL) and this solution was diluted with 65% acetonitrile/35% 0.1% phosphoric acid in water (250 mL). A 75 mL aliquot was injected onto a preparative HPLC column. The column consisted of an Amicon Matrex LC ODS Silica (10 cm ID x 50 cm) with 100Å 20 µm particle size. Elution was carried out with a mobile phase of 65% CH<sub>3</sub>CN/35% 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O at a flow rate of 250 mL/min, collecting fractions at four minute intervals to provide 1 L fractions, and detecting at 220 nm. The fractions were pooled as follows: fractions #3-4; fractions #5-6; fraction #7; fractions #8-9; fractions #10-11; fractions #12-14; fractions #15-17 and fractions #18-20. A total of 10 injections were performed, combining pooled fractions from each run, to give Cut 3-4, Cut 5-6, Cut 7, Cut 8-9, Cut 10-11, Cut 12-14, Cut 15-17, Cut 18-20. Cuts 5-6, 8-9, 12-14, and 18-20 were further fractionated using preparative HPLC.

20

Cut 5-6:

Cut 5-6 was partitioned with an equal volume of EtOAc. The organic layer was evaporated to dryness. One half of the dry residue from Cut 5-6 was dissolved in 3 mL of CH<sub>3</sub>CN and 1 mL of 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O was added. This solution was injected on a preparative HPLC column (Dynamax C8 8 µm 41.4 mm ID x 250 mm L + guard) eluting with 60% CH<sub>3</sub>CN/40% 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O at a flow rate of 24 mL/min. Fractions were collected at 1 minute intervals (detecting wavelength = 220 nm). Each of the following fractions was analyzed by analytical HPLC and then extracted with an equal volume of EtOAc. The organic layer was evaporated to dryness to afford dry material.

30

The analytical HPLC conditions involved a reverse-phase silica-based HPLC column (Dynamax C8, 60 Angstroms, 8 $\mu$ m particles, 4.6 mm ID x 250 mm with guard column) eluting at room temperature at a flow rate of 1 ml/min with a mobile phase of 55% CH<sub>3</sub>CN/45% of 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O. The detection wavelength was 205 nm. Relative retention time to zaragozic acid C (RRT) is calculated for each compound as the ratio of the retention time of a given major peak divided by the retention time of zaragozic acid C (30.5 min.).

Fraction 27 (6.6 mg) and 28 (5.1 mg): Each contain at least 2 components. The components have the molecular weights 710 and 770 by FAB-MS (observed [M-H]<sup>-</sup> at m/z 709 and 769, respectively). The 710 component has the molecular formula C<sub>38</sub>H<sub>46</sub>O<sub>13</sub> as determined by HR-EI measurement of the trimethyl ester derivatives (calc. for C<sub>38</sub>H<sub>46</sub>O<sub>13</sub> + (CH<sub>2</sub>)<sub>3</sub> -H<sub>2</sub>O; 734.3302 found 734.3299). Relative Retention Time (RRT) = 0.30.

Fraction 30 (7 mg): contains a major component. The component has the molecular weight 728 by FAB-MS (observed [M-H]<sup>-</sup> at m/z 727. It has the molecular formula C<sub>39</sub>H<sub>52</sub>O<sub>13</sub>, as determined by HR-EI measurement of the trimethyl ester derivatives (calc. for C<sub>39</sub>H<sub>52</sub>O<sub>13</sub> + (CH<sub>2</sub>)<sub>3</sub> -H<sub>2</sub>O; 752.3772, found 752.3750). Relative Retention Time (RRT) = 0.37

Fraction 31 (21.1 mg): contains at least 2 major components. The components have the molecular weights 698 and 712 by FAB-MS (observed [M-H]<sup>-</sup> at m/z 697 and 711 respectively). They have the molecular formulas C<sub>37</sub>H<sub>46</sub>O<sub>13</sub> and C<sub>38</sub>H<sub>48</sub>O<sub>13</sub> as determined by HR-EI

measurement of the trimethyl ester derivatives (calc. for  $C_{37}H_{46}O_{13} + (CH_2)_3 - H_2O$ ; 722.3233, found 722.3268 and calc for  $C_{38}H_{48}O_{13} + (CH_2)_3 - CH_3COOH$ ; 736.3459 found 736.3444).

5 Relative Retention Time (RRT) = 0.38

Fraction 33 (8.1 mg): contains at least 2 components. The components have the molecular weight 710 and 768 by FAB-MS (observed  $[M-H]^-$  at  $m/z$  709 and 767


10 respectively). The 710 component has the molecular formula  $C_{38}H_{46}O_{13}$  as determined by HR-EI measurement of the trimethyl ester derivative (calc. for  $C_{38}H_{46}O_{13} + (CH_2)_3 - H_2O$ ; 734.3302, found 734.3291. Relative Retention Time (RRT) = 0.41.

15

Fraction 36 (10.1 mg): The major component in this fraction has a molecular weight of 700 by FAB-MS (observed  $[M-H]^-$  at  $m/z$  699). The molecular formula  $C_{37}H_{48}O_{13}$  was determined by HR-EI measurement of the trimethylester derivative [calc. for  $(C_{37}H_{48}O_{13} + (CH_2)_3 - H_2O)$  724.3459, found 724.3458]. The major component has a RRT = 0.46 and may be 4'-O-desacetyl-4'',5''-dihydro-6''-desmethylzaragozic acid C, MW 700, of the structure:

25

30




Fraction 38 (31.2 mg): The major component in this fraction has a molecular weight of 712 by FAB-MS (observed  $[M-H]^-$  at  $m/z$  711). The molecular formula  $C_{38}H_{48}O_{13}$  was determined by HR-EI measurement of the trimethylester derivative [calc. for  $(C_{38}H_{48}O_{13} + (CH_2)_3 - H_2O)$  736.3459, found 736.3451]. The major component has RRT = 0.49 and may be 4'-O-desacetyl-zaragozic acid C MW 712.

10 Cut 7:

Contains 4'-O-desacetyl-zaragozic acid C MW 712 (10.0 g), RRT = 0.49

15



20

Cut 8-9:

One tenth of Cut 8-9 was partitioned with an equal volume of EtOAc. The organic layer was evaporated to dryness. The dry residue was dissolved in 2 mL of  $CH_3CN$ /0.5 mL of 0.1%  $H_3PO_4$  in  $H_2O$ . This solution was injected on a preparative HPLC column (Dynamax C8 8 $\mu$ m 41.4 mm ID x 250 mm L + guard) eluting with 65%  $CH_3CN$ /35% of 0.1%  $H_3PO_4$  in  $H_2O$  at a flow rate of 24 mL/min. Fractions were collected at 1 minute intervals (detecting wavelength = 220 nm). Fractions were combined as shown below and each of the pools was analyzed by analytical HPLC and then extracted with an


25

30

equal volume of EtOAc. The organic layer was evaporated to dryness to afford dry material.

The analytical HPLC conditions involved a reverse-phase silica-based HPLC column (Dynamax C8, 60 Angstroms, 8  $\mu$ m particles, 4.6 mm ID x 250 mm with guard column) eluting at room temperature at a flow rate of 1 ml/min with a mobile phase of 55% CH<sub>3</sub>CN/45% of 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O. The detection wavelength was 205 nm. Relative retention time to zaragozic acid C (RRT) is calculated for each compound as the ratio of the retention time of a given major peak divided by the retention time of zaragozic acid C (30.5 min).

Fractions 31-32: contains 2 compounds MW 714, 738. The 714 component has a molecular weight of 714 by FAB-MS (observed [M-H]<sup>-</sup> at m/z 713). Its molecular formula C<sub>38</sub>H<sub>50</sub>O<sub>13</sub> was determined by HR-EI measurement of the trimethylester derivative [calc. for (C<sub>38</sub>H<sub>50</sub>O<sub>13</sub> + (CH<sub>2</sub>)<sub>3</sub> -H<sub>2</sub>O) 738.3615, found 738.3606]. The 714 component has RRT = 0.58 and may be 4'-O-desacetyl-4'',5''-dihydro-zaragozic acid C, of the structure:




30 The 738 component has a molecular weight of 738 by FAB-MS (observed [M-H]<sup>-</sup> at m/z 737) and a molecular formula C<sub>40</sub>H<sub>50</sub>O<sub>13</sub> as determined by HR-EI



measurement of the trimethylester derivative [calc. for  $(C_{40}H_{50}O_{13} + (CH_2)_3 - H_2O)$  762.3615, found 762.3612].  
RRT = 0.58.

- 5 Fractions 36-37: contains at least 2 compounds MW 726  
and MW 740. The 726 component has a molecular weight  
of 726 by FAB-MS (observed  $[M-H]^-$  at  $m/z$  725). The  
molecular formula  $C_{39}H_{50}O_{13}$  was determined by HR-EI  
measurement of the dimethylester derivative [calc. for  
10  $(C_{39}H_{50}O_{13} + (CH_2)_2 - H_2O)$  736.3459, found 736.3458].  
It may be 4'-O-desacetyl-4'',5''-dihydro-zaragozic acid C  
monomethyl ester. RRT = 0.74.

15



20

- The second component has a molecular weight of 740 by  
FAB-MS (observed  $[M-H]^-$  at  $m/z$  739). The molecular  
formula  $C_{39}H_{48}O_{14}$  was determined by HR-EI measurement  
of the trimethylester derivative [calc. for  $(C_{39}H_{48}O_{14}$   
25  $+ C_3H_6 - CH_3COOH)$  722.3302, found 722.3304].

- Fractions 38-39: This contains a compound of MW 752,  
by FAB-MS (observed  $[M-H]^-$  at  $m/z$  751). The molecular  
formula  $C_{40}H_{48}O_{14}$  was determined by HR-EI measurement  
30 of the trimethylester derivative [calc. for  $(C_{40}H_{48}O_{14}$   
 $+ C_3H_6 - CH_3COOH)$  734.3302, found 734.3299]. RRT =  
0.81. It may be 8'',9''-dehydro-zaragozic acid C, of the  
structure:



10

**RRT = 1.00.**




20

Two thirds of Cut 12-14 was partitioned with an equal volume of EtOAc. The organic layer was evaporated to dryness. One half of the dry residue was dissolved in 3 mL of CH<sub>3</sub>CN/1 mL of 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O. This solution was injected on a preparative HPLC column (Dynamax C8 8µm 41.4 mm ID x 250 mm L + guard) eluting with 70% CH<sub>3</sub>CN/30% of 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O at a flow rate of 24 mL/min. Fractions were collected at 1 min intervals (detecting wavelength = 220 nm). Fractions were combined as shown below and each of the pools was analyzed by analytical HPLC and then extracted with an equal volume of EtOAc. The organic layer was evaporated to dryness to afford dry material.

The analytical HPLC conditions involved a reverse-phase silica-based HPLC column (Dynamax C8, 60 Angstroms, 8  $\mu$ m particles, 4.6 mm ID x 250 mm with guard column) eluting at room temperature at a flow rate of 1 ml/min with a mobile phase of 65% CH<sub>3</sub>CN/35% of 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O. The detection wavelength was 205 nm. Relative retention time to zaragozic acid C (RRT) is calculated for each compound as the ratio of the retention time of a given major peak divided by the retention time of zaragozic acid C. (11.5 min).

Fractions 30-31 (284.5 mg): contains zaragozic acid C MW 754

Fractions 35-36 (78.9 mg): contains a compound of MW 756 by FAB-MS (observed [M-H]<sup>-</sup> at m/z 755). The molecular formula C<sub>40</sub>H<sub>52</sub>O<sub>14</sub> was determined by HR-EI measurement of the trimethylester derivative [calc. for (C<sub>40</sub>H<sub>52</sub>O<sub>14</sub> + (CH<sub>2</sub>)<sub>3</sub> -CH<sub>3</sub>COOH) 738.2615, found 738.3611]. RRT = 1.10. It may be 4'',5''-dihydro-zaragozic acid C:




30 Fractions 39-40 (34.6 mg): contains 4'-desacetoxy zaragozic acid C MW 696

Cut 15-17:

Contains 4'-desacetoxy zaragozic acid C (L-735,021); MW 696 (9.7 g). RRT = 1.3.

5

10

Cut 18-20:

15

20

25

Half of cut 18-20 was partitioned with an equal volume of EtOAc. The organic layer was evaporated to dryness. One fifth of the dry residue was dissolved in 3 mL of CH<sub>3</sub>CN/1 mL of 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O. This solution was injected on a preparative HPLC column (Dynamax C8 8  $\mu$ m 41.4 mm ID x 250 mm L + guard) eluting with 75% CH<sub>3</sub>CN/25% 0.1% H<sub>3</sub>PO<sub>4</sub> in H<sub>2</sub>O at a flow rate of 24 mL/min. Fractions were collected at 1 min intervals (detecting wavelength = 220 nm). Fractions were combined as shown below and each of the pools was analyzed by analytical HPLC and then extracted with an equal volume of EtOAc. The organic layer was evaporated to dryness to afford dry material.


30

The analytical HPLC conditions involved a reverse-phase silica-based HPLC column (Dynamax C8, 60 Angstroms, 8  $\mu$ m particles, 4.6 mm ID x 250 mm with guard column) eluting at room temperature at a flow rate of 1 mL/min with a mobile phase of 65% CH<sub>3</sub>CN/35%

of 0.1%  $\text{H}_3\text{PO}_4$  in  $\text{H}_2\text{O}$ . The detection wavelength was 205 nm. Relative retention time to zaragozic acid C (RRT) is calculated for each compound as the ratio of the retention time of a given major peak divided by the retention time of zaragozic acid C. (11.5 min).

Fractions 31-32 (34.6 mg): contains 4'-desacetoxy zaragozic acid C MW 696. RRT = 1.3.

Fractions 35-36 (19.4 mg): The major component in this fraction has a molecular weight of 698 by FAB-MS (observed  $[\text{M}-\text{H}]^-$  at  $m/z$  697). The molecular formula  $\text{C}_{38}\text{H}_{50}\text{O}_{12}$  was determined by HR-EI measurement of the trimethylester derivative [calc. for  $(\text{C}_{38}\text{H}_{50}\text{O}_{12} + (\text{CH}_2)_3$ ) 740.3771, found 740.3770]. May contain 4'-desacetoxy dihydrozaragozic acid C MW 698. RRT = 1.5.



25

Fractions 46-47 (9.1 mg): contains a major component with the molecular weight of 696 by FAB-MS (observed  $[\text{M}-\text{H}]^-$  at  $m/z$  695). The molecular formula  $\text{C}_{38}\text{H}_{48}\text{O}_{12}$  was determined by HR-EI measurement of the trimethylester derivative [calc. for  $(\text{C}_{38}\text{H}_{48}\text{O}_{12} + (\text{CH}_2)_3$ ) 738.3615, found 738.3618]. RRT = 2.0.

- Fractions 52-54 (15.8 mg): contain two components MW 676 and 736. The 676 component, FAB-MS (observed  $[M-H]^-$  at  $m/z$  675), forms a trimethylester observed at  $m/z$  700 ( $676 + (CH_2)_3 - H_2O$ ). The 736 component
- 5 [FAB-MS (observed  $[M-H]^-$  at  $m/z$  735)] has a molecular formula  $C_{40}H_{48}O_{13}$  as determined by HR-EI measurement of the dimethylester derivative [calc. for  $(C_{40}H_{48}O_{13} + C_2H_4 - CH_3COOH)$  718.3353, found 718.3354]. RRT = 2.5.
- 10 Fractions 61-62 (9.7 mg): contains a major component MW 678, by FAB-MS (observed  $[M-H]^-$  at  $m/z$  677) which forms a trimethylester at  $m/z$  720 ( $678 + (CH_2)_3$ ). RRT = 2.7.
- 15 Fractions 65-66 (11.2 mg): contains a major component MW 750, by FAB-MS (observed  $[M-H]^-$  at  $m/z$  749). RRT = 2.8.

20 The isolated compounds of this Example are summarized in the following Table. It will be understood that the particulars of this Table are subject to minor revisions as additional tests become available subsequent to the filing of this application.

25

30

TABLE

	<u>MW</u>	<u>Molecular Formula</u>	<u>Relative Retention Time (RRT)</u>	<u>Cut/Fractions*</u>
5	676	n/a	2.5	18-20/52-54
	678	n/a	2.7	18-20/61-62
	696	C <sub>38</sub> H <sub>48</sub> O <sub>12</sub>	1.3	15-17
	696	C <sub>38</sub> H <sub>48</sub> O <sub>12</sub>	2.0	18-20/46-47
	698	C <sub>37</sub> H <sub>46</sub> O <sub>13</sub>	0.38	5-6/31
10	698	C <sub>38</sub> H <sub>50</sub> O <sub>12</sub>	1.5	18-20/35-36
	700	C <sub>37</sub> H <sub>48</sub> O <sub>13</sub>	0.46	5-6/36
	710	C <sub>38</sub> H <sub>46</sub> O <sub>13</sub>	0.30	5-6/27-28
	710	C <sub>38</sub> H <sub>46</sub> O <sub>13</sub>	0.41	5-6/33
	712	C <sub>38</sub> H <sub>48</sub> O <sub>13</sub>	0.38	5-6/31
15	712	C <sub>38</sub> H <sub>48</sub> O <sub>13</sub>	0.49	7,5-6/38
	714	C <sub>38</sub> H <sub>50</sub> O <sub>13</sub>	0.58	8-9/31-32
	726	C <sub>39</sub> H <sub>50</sub> O <sub>13</sub>	0.74	8-9/36-37
	728	C <sub>39</sub> H <sub>52</sub> O <sub>13</sub>	0.37	5-6/30
	736	C <sub>40</sub> H <sub>48</sub> O <sub>13</sub>	2.5	18-20/52-54
20	738	C <sub>40</sub> H <sub>50</sub> O <sub>13</sub>	0.58	8-9/31-32
	740	C <sub>39</sub> H <sub>48</sub> O <sub>14</sub>	0.74	8-9/36-37
	750	n/a	2.8	18-20/65-66
	752	C <sub>40</sub> H <sub>48</sub> O <sub>14</sub>	0.81	8-9/38-39
	754	C <sub>40</sub> H <sub>50</sub> O <sub>14</sub>	1.00	10,11
25	756	C <sub>40</sub> H <sub>52</sub> O <sub>14</sub>	1.10	12-14/35-36
	768	n/a	0.41	5-6/33
	770	n/a	0.30	5-6/27-28

\* Cut refers to combined fractions from the large scale HPLC /  
 30 fractions refers to HPLC fractions from the fractionation of  
 the cuts.

EXAMPLE 4Preparation of an Ammonium Salt

5 A 0.1 mmol sample of the free acid of a  
compound of formula (I) is dissolved in 10 ml of ethyl  
acetate. The resulting solution is saturated with  
gaseous ammonia upon which the ammonium salt  
precipitates from solution.

10

EXAMPLE 5Preparation of a Potassium Salt

15 A solution of 0.1 mmol of the free acid of a  
compound of formula (I) in 10 ml of methanol is treated  
with an aqueous or methanolic solution containing 0.3  
mmol of potassium hydroxide. Evaporation of the  
solvent affords the tri-potassium salt. Addition of  
between 0.1 and 0.3 mmol of potassium hydroxide yields  
analogously mixtures of the mono-potassium,  
20 di-potassium and tri-potassium salts whose composition  
depends upon the exact amount of potassium hydroxide  
added.

In a similar fashion the sodium and lithium  
salts can be formed.

25

EXAMPLE 6Preparation of a Calcium Salt

30 A solution of 0.1 mmol of the free acid of a  
compound of formula (I) in 20 ml of 6:4 methanol/ water  
is treated with an aqueous solution of 0.1 mmol of  
calcium hydroxide. The solvents are evaporated to give  
the corresponding calcium salt.



EXAMPLE 7Preparation of an Ethylenediamine Salt

5 A solution of 0.1 mmol of the free acid of a compound of formula (I) in 10 ml of methanol is treated with 0.1 mmol of ethylenediamine. Evaporation of the solvent affords the ethylenediamine salt. The procedure can also be applied to the preparation of the N,N''-dibenzylethylenediamine salt.

10

EXAMPLE 8Preparation of a Tris(hydroxymethyl)aminomethane Salt

15 To a solution of 0.1 mmol of the free acid of a compound of formula (I) in 10 ml of methanol is added from 0.1 to 0.3 mmol of tris(hydroxymethyl)aminomethane dissolved in 10 ml of methanol. Evaporation of the solvent gives a corresponding salt form, the exact composition of which is determined by the molar ratio of amine added. Similarly prepared are the salts of L-ornithine, L-lysine, and N-methylgluacamine.

20

EXAMPLE 9

25 Preparation of an L-arginine Salt

A solution of 0.1 mmol of the free acid of a compound of formula (I) in 20 ml of 6:4 methanol/ water is treated with an aqueous solution of 0.1 to 0.3 mmol of L-arginine. Evaporation of the solvent affords the title salt, the exact composition of which is determined by the molar ratio of amino acid to the free acid of formula (I) used.

30

Similarly prepared are the salts of L-ornithine, L-lysine and N-methylglucamine.

EXAMPLE 10

5                   Preparation of A trimethyl ester


A solution of 2 mg of a free acid of compound of formula (I) in 0.5 ml of acetonitrile is treated at room temperature with 10 equivalents of DBU and 10  
10 equivalents of MeI. After 2 hours the reaction is diluted with 10 ml of dichloromethane and washed successively with 10 ml of 0.1 M phosphoric acid, 10 ml of water; 10 ml of saturated sodium bicarbonate and 10 ml of water. After drying over sodium sulfate, the  
15 organic layer is concentrated and the residue is chromatographed on silica gel using mixtures of hexane and ethyl acetate to give a trimethyl ester

The method of Example 7 is also suitable for  
20 the preparation of other ester derivatives such as 1) ethyl and other lower alkyl esters and 2) benzyl and substituted benzyl esters.

While the foregoing specification teaches the  
25 principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations, modifications, deletions or additions of procedures and  
30 protocols described herein, as come within the scope of the following claims and its equivalents.




WHAT IS CLAIMED IS:

1. A compound of the structural formula (I)




or pharmaceutically acceptable salt thereof,  
15 wherein

$R^1$  is selected from the group consisting of




$R^2$  is selected from the group consisting of


5




10



15



, or



20  $Z^1$ ,  $Z^2$  and  $Z^3$  are each independently selected from;

a) H;

b)  $C_{1-5}$ alkyl;


c)  $C_{1-5}$ alkyl substituted with  
a member of the group consisting of:

25

i) phenyl,

ii) phenyl substituted with methyl,  
methoxy, halogen (Cl, Br, I, F) or  
hydroxy;


30 with the proviso that when  $R^2$  is



5.


then R<sup>1</sup> is neither

10



nor,

15




2. A compound according to Claim 1, wherein Z<sup>1</sup>, Z<sup>2</sup> and Z<sup>3</sup> are each independently H or CH<sub>3</sub>.

20

3. A compound according to Claim 1, of the structure


25



30


4. A compound according to Claim 1, of the structure

5



5. A compound according to Claim 1, of the structure


10



15

6. A compound according to Claim 1, of the structure


20




25

7. A compound according to Claim 1, of the structure


30



8. A compound according to Claim 1, of the structure



9. A compound according to Claim 1, of the structure



10. A compound with a molecular weight of 676, and a relative retention time of 2.5 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 65%  $\text{CH}_3\text{CN}$ /about 35% of about 0.1%  $\text{H}_3\text{PO}_4$  in  $\text{H}_2\text{O}$ .

11. A compound with a molecular weight of 678, and a relative retention time of 2.7 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 65%  $\text{CH}_3\text{CN}$ /about 35% of about 0.1%  $\text{H}_3\text{PO}_4$  in  $\text{H}_2\text{O}$ .

12. A compound with a molecular weight of 696, a molecular formula of  $C_{38}H_{48}O_{12}$ , and a relative retention time of 2.0 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room  
5 temperature at flow rate of about 1 ml/minute with a mobile phase of about 65%  $CH_3CN$ /about 35% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

13. A compound with a molecular weight of 698, a molecular formula of  $C_{37}H_{46}O_{13}$ , and a relative  
10 retention time of 0.38 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  
15  $H_3PO_4$  in  $H_2O$ .

14. A compound with a molecular weight of 698, a molecular formula of  $C_{38}H_{50}O_{12}$ , and a relative  
retention time of 1.5 on a reverse-phase silica-based  
20 HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 65%  $CH_3CN$ /about 35% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

25 15. A compound with a molecular weight of 700, a molecular formula of  $C_{37}H_{48}O_{13}$ , and a relative retention time of 0.46 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room  
temperature at flow rate of about 1 ml/minute with a  
30 mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .



16. A compound with a molecular weight of 710, a molecular formula of  $C_{38}H_{46}O_{13}$ , and a relative retention time of 0.30 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room  
5 temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

17. A compound with a molecular weight of 710, a molecular formula of  $C_{38}H_{46}O_{13}$ , and a relative  
10 retention time of 0.41 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  
15  $H_3PO_4$  in  $H_2O$ .

18. A compound with a molecular weight of 712, a molecular formula of  $C_{38}H_{48}O_{13}$ , and a relative  
20 retention time of 0.38 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  
 $H_3PO_4$  in  $H_2O$ .

19. A compound with a molecular weight of 712, a molecular formula of  $C_{38}H_{48}O_{13}$ , and a relative  
25 retention time of 0.49 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a  
30 mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

20. A compound with a molecular weight of 714, a molecular formula of  $C_{38}H_{50}O_{13}$ , and a relative retention time of 0.58 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room  
5 temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

21. A compound with a molecular weight of 726, a molecular formula of  $C_{39}H_{50}O_{13}$ , and a relative  
10 retention time of 0.74 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  
15  $H_3PO_4$  in  $H_2O$ .

22. A compound with a molecular weight of 728, a molecular formula of  $C_{39}H_{52}O_{13}$ , and a relative retention time of 0.37 on a reverse-phase silica-based  
20 HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

23. A compound with a molecular weight of 736, a molecular formula of  $C_{40}H_{48}O_{13}$ , and a relative retention time of 2.5 on a reverse-phase silica-based  
25 HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 65%  $CH_3CN$ /about 35% of about 0.1%  
30  $H_3PO_4$  in  $H_2O$ .

24. A compound with a molecular weight of 738, a molecular formula of  $C_{40}H_{50}O_{13}$ , and a relative retention time of 0.58 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

25. A compound with a molecular weight of 740, a molecular formula of  $C_{39}H_{48}O_{14}$ , and a relative retention time of 0.74 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at a flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

26. A compound with a molecular weight of 750, and a relative retention time of 2.8 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 65%  $CH_3CN$ /about 35% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

27. A compound with a molecular weight of 752, a molecular formula of  $C_{40}H_{48}O_{14}$ , and a relative retention time of 0.81 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

28. A compound with a molecular weight of 756, a molecular formula of  $C_{40}H_{52}O_{14}$ , and a relative retention time of 1.10 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room  
5 temperature at flow rate of about 1 ml/minute with a mobile phase of about 65%  $CH_3CN$ /about 35% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

29. A compound with a molecular weight of 768, and a relative retention time of 0.41 on a  
10 reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .  
15

30. A compound with a molecular weight of 770, and a relative retention time of 0.30 on a reverse-phase silica-based HPLC column (Dynamax C8) eluting at about room temperature at flow rate of about  
20 1 ml/minute with a mobile phase of about 55%  $CH_3CN$ /about 45% of about 0.1%  $H_3PO_4$  in  $H_2O$ .

31. A pharmaceutical composition comprising a nontoxic therapeutically effective amount of a  
25 compound of Claim 1 and a pharmaceutically acceptable carrier.

32. A pharmaceutical composition comprising  
a nontoxic therapeutically effective amount of a  
compound of Claim 1 in combination with a  
pharmaceutically acceptable nontoxic cationic polymer  
5 capable of binding bile acids in a non-reabsorbable  
form in the gastrointestinal tract and pharmaceutically  
acceptable carrier.

33. A pharmaceutical composition comprising  
10 a nontoxic therapeutically effective amount of a  
compound of Claim 1 in combination with a nontoxic  
therapeutically effective amount of a cholesterol  
lowering agent selected from the group consisting of:

- 15 (a) HMG-CoA reductase inhibitor;  
(b) HMGCoA synthase inhibitor;  
(c) Squalene epoxidase inhibitor;  
(d) Probucol;  
(e) Niacin;  
20 (f) Gemfibrozil;  
(g) Clofibrate.

34. A composition of Claim 33 wherein the  
composition comprises a compound of Claim 1 and an  
25 HMG-CoA reductase inhibitor.

35. A composition of Claim 34 wherein the  
HMG-CoA reductase inhibitor is selected from  
lovastatin, simvastatin, pravastatin and fluvastatin.  
30

36. The use of a compound as claimed in claim 1 for the manufacture of a medicament for treating hypercholesterolemia.

5

37. The use of a compound as claimed in claim 1 for the manufacture of a medicament for inhibiting squalene synthetase.

10

38. A method for inhibiting fungal growth comprising applying to the area where growth is to be controlled an anti-fungally effective amount of a compound of Claim 1.

15

39. The use of a compound as claimed in claim 1 for the manufacture of a medicament for treating cancer.

20

40. The use of a compound as claimed in claim 1 for the manufacture of a medicament for inhibiting farnesyl-protein transferase and farnesylation of the oncogene protein Ras.

25

41. A method of making a compound as in any of Claims 1-30, comprising the steps of:

30

- (a) growing a sample of MF5465, to give a fermentation broth;
- (b) extracting the fermentation broth; and
- (c) chromatographing to give the compound.

- 52 -

**Patents Act 1977**  
**Examiner's report to the Comptroller under Section 17**  
**(The Search report)**

Application number  
 GB 9317770.7

**Relevant Technical Fields**

(i) UK Cl (Ed.L)      C2C CBC

(ii) Int Cl (Ed.5)      C07D

Search Examiner  
 D S LUCAS

Date of completion of Search  
 16 NOVEMBER 1993

**Databases (see below)**

(i) UK Patent Office collections of GB, EP, WO and US patent specifications.

(ii) ONLINE DATABASE: CAS ONLINE

Documents considered relevant following a search in respect of Claims :-  
 1-41

**Categories of documents**

- |  |   |
|--|---|
| <p><b>X:</b> Document indicating lack of novelty or of inventive step.</p> <p><b>Y:</b> Document indicating lack of inventive step if combined with one or more other documents of the same category.</p> <p><b>A:</b> Document indicating technological background and/or state of the art.</p> | <p><b>P:</b> Document published on or after the declared priority date but before the filing date of the present application.</p> <p><b>E:</b> Patent document published on or after, but with priority date earlier than, the filing date of the present application.</p> <p><b>&amp;:</b> Member of the same patent family; corresponding document.</p> |
|--|---|

Category	Identity of document and relevant passages	Relevant to claim(s)
X	EP 0475706 A1 (MERCK) see Claims 1-10	1-41
X	US 5026554 A (MERCK) see Claims 1 and 6 and lines 23-35 in column 3	1-41

**Databases:** The UK Patent Office database comprises classified collections of GB, EP, WO and US patent specifications as outlined periodically in the Official Journal (Patents). The on-line databases considered for search are also listed periodically in the Official Journal (Patents).